CONTENTS

Influence of the spectral absorption curves for nitrogen-bearing compounds based on a computer library
L.A. Gribov, A.T. Todorovsky and S.Ya. Plotkin (Moscow, U.S.S.R.) 1
Influence de la structure des cétones et de l'acidité des phénols sur leur mode de complexation. Partie I. Comportement global et échelles d'acidité et de basicité
A. Massat, A. Cossé-Barbi and J.P. Doucet (Paris, France) 13
Comparison of theoretical and experimental ionization potentials of nicotine and related molecules
F.W. Kuhn, D. Lilly-Leister, J. Kao and A.C. Lilly (Richmond, VA, U.S.A.) 37
Calculations of structures of styrenes and stilbenes by molecular mechanics
S. Tsuzuki, K. Tanabe, Y. Nagawa and H. Nakanishi (Tsukuba, Ibaraki, Japan) 45
Ab initio studies on proton transfer involving Schiff base and related nitrogen compounds
N. Sreerama and S. Vishveshwara (Bangalore, India) 53
A molecular mechanics force field for conformational analysis of simple thiocarbonyl compounds
R. Fausto, J.J.C. Teixeira-Dias (Coimbra, Portugal) and P.R. Carey (Ottawa, Ont., Canada) 61
Molecular structure of cyclododecane, $C_{12}H_{24}$, as determined by electron diffraction and molecular mechanics
E.G. Atavin, V.S. Mastryukov (Moscow, U.S.S.R.), N.L. Allinger (Athens, GA, U.S.A.), A. Almenningen and R. Seip (Oslo, Norway) 87
On the zwitterionic nature of strongly twisted push-pull ethylenes: the case of 3-(1,3-diisopropyl-2-imidazolidinylidene)-2,4-pentanedione monohydrate, studied by X-ray diffraction and ab initio calculations
R. Destro, U. Cosentino, G. Moro, E. Ortoleva and T. Pilati (Milan, Italy) 97
Preferred conformations of methylthio-acetone and -acetophenone, methylsulphonyl-acetone and -acetophenone, from molecular-mechanics and dipole-moment techniques
H. Lumbroso, D.M. Bertin (Paris, France), P.R. Olivato, É. Bonfada, M.G. Mondino (São Paulo, Brazil) and Y. Hase (Campinas, Brazil) 113
A conformational study of oligopeptides containing gly–pro sequence in the solid state by ^{13}C CP-MAS NMR
S. Ando, K. Matsumoto, I. Ando (Tokyo, Japan), A. Shoji and T. Ozaki (Gunma, Japan) 123
Zero-point average structure of methanol
T. Iijima (Tokyo, Japan) 137
Molecular structure and conformation of *trans*-1,3-dichloro-1-propene as determined by gas-phase electron diffraction
Q. Shen and C. Nunn (Hamilton, NY, U.S.A.) 143
The molecular structure and conformation of *trans*-1,2,3-trichloropropene as determined by gas-phase electron diffraction
Q. Shen (Hamilton, NY, U.S.A.) 151
Raman spectra and rotational isomerism of dimethylphosphorochloridate and trimethyl phosphate
K. Taga, N. Hirabayashi, T. Yoshida and H. Okabayashi (Nagoya, Japan) 157
Infrared and Raman spectra, vibrational assignment, and conformational stability of methacryloyl bromide
J.R. Durig, W. Zhao, R.J. Berry and T.S. Little (Columbia, SC, U.S.A.) 169
Conformational stability, barriers to internal rotation, normal coordinate analyses, and vibrational assignments of bromoacetyl halides
J.R. Durig, H.V. Phan and T.S. Little (Columbia, SC, U.S.A.) 187
Resonance Raman and absorption studies of the configurations of photochromic 3-alkyl-substituted 1,5-diphenylformazans: steric effect of the substituent
H. Hiura and H. Takahashi (Tokyo, Japan) 221

Study of the configurations of 3-aryl-substituted 1,5-diphenylformazans by resonance Raman and absorption spectroscopy: steric and conjugation effects of the substituent
H. Hiura and H. Takahashi (Tokyo, Japan) 235
Change of the proton potential from a single to a double minimum in intramolecular hydrogen bonds with increasing pK_a of the phenolic group in 4-R-Mannich base *N*-oxides
B. Brzezinski, J. Olejnik (Poznan, Poland), G. Zundel and R. Krämer (Munich, F.R.G.) 247
Spectroscopic properties of $Cr(CN)_6^{3-}$ ion trapped in NaCl, KCl, RbCl and KBr hosts
J. Hanuza, K. Hermanowicz, B. Jeżowska-Trzebiatowska (Wroclaw, Poland) and S. Waplak (Poznań, Poland) 255
The structure investigation of a new product formed in the course of nitration of apovincaminic acid ethyl ester by NMR methods
G. Tóth, Cs. Szántay Jr., I. Moldvai, Cs. Szántay (Budapest, Hungary), W. Dietrich and A. Vedres (Bochum, F.R.G.) 271
Crystal and molecular structure of the two MACRO-ANSA and MACRO-SPIRO isomeric cyclotriphosphazenes $N_3P_3Cl_4$[HN–$(CH_2)_3$–O–$(CH_2)_4$–O–$(CH_2)_3$–NH]
T.S. Cameron, A. Linden (Halifax, NS, Canada), A. El Bakili, P. Castera, J.-P. Faucher, M. Graffeuil, F. Sournies and J.-F. Labarre (Toulouse, France) 281
Crystal and molecular structure of the SPIRANSA and DISPIRANSA cyclophosphazenic derivatives from spermidine and spermine, $N_3P_3Cl_3$[HN–$(CH_2)_3$–N–$(CH_2)_4$–NH] and $N_3P_3Cl_2$[HN–$(CH_2)_3$–N–$(CH_2)_4$–N–$(CH_2)_3$N–NH]
T.S. Cameron, A. Linden (Halifax, NS, Canada), G. Guerch, J.-P. Bonnet and J.-F. Labarre (Toulouse, France) 295
Crystal and molecular structure of a new allotropic variety of the MEGA-SPIRO cyclophosphazene from diethylene glycol bis-(3-aminopropyl)-ether, $N_3P_3Cl_4$-[HN–$(CH_2)_3$–O–$(CH_2)_2$–O–$(CH_2)_3$–$(CH_2)_3$–$(CH_2)_3$–NH]
J. Jaud, F. Sournies and J.-F. Labarre (Toulouse, France) 305
Comparison between optimized and crystal structure of R5135, a potent steroidic GABA-A antagonist: ab initio molecular orbital charge population analysis
T. Boulanger, D.P. Vercauteren, G. Evrard and F. Durant (Namur, Belgium) 315

Short communication
On the *T* symmetry of the molecule tetrakis(trimethylsilyl) methane in the gas phase
B. Beagley, R.G. Pritchard and J.O. Titiloye (Manchester, Gt. Britain) 323

Book reviews 325
Author index 335
Subject index 337